

# THE IDEAL GIFT FOR EVERY FAMILY!

## BIRD BOX CAMERA



*SEE & HEAR FROM INSIDE YOUR BIRD BOX DAY & NIGHT*

Quick and easy to fit into Bird Box or other weatherproof location

No tuning required - view at the touch of a button

Just one cable to connect to TV or Video & 12V adaptor

Infrared LED's do not disturb wildlife

### Package Includes

1 X Bird Box Monochrome Camera with Audio & IR LED's



1 x 30m Connection Cable, Connect From Camera to TV

1 x SCART Plug

1 x Power Adaptor for Camera

**Nature Programmes such as BBCs "Spring Watch" like this product**

Our DIY & Garden Gold Award Winning Cameras proved popular as a great gift for the notoriously difficult-to-buy-for market of Fathers & Grandfathers.

The Bird Box Camera is seen as an educational and fun product that can give pleasure to young and old alike.



# When to feed wild birds



Although winter feeding benefits birds most, food shortages can occur at any time of the year. By feeding the birds year round, you'll give them a better chance to survive the periods of food shortage whenever they may occur.

## **Autumn and winter**

At this time of year, put out food and water on a regular basis. In severe weather, feed twice daily if you can: in the morning and in the early afternoon.

Birds require high energy (high fat) foods during the cold winter weather to maintain their fat reserves to survive the frosty nights. Use only good quality food and scraps.

Always adjust the quantity given to the demand, and never allow uneaten foods to accumulate around the feeders. Once you establish a feeding routine, try not to change it as the birds will become used to it and time their visits to your garden accordingly.

## **Spring and summer**

Only feed selected foods at this time of year. Good hygiene is vital, or feeding may do more harm than good.

During the summer months, birds require high protein foods, especially while they are moulting.

Black sunflower seeds, pinhead oatmeal, soaked sultanas, raisins and currants, mild grated cheese, mealworms, waxworms, mixes for insectivorous birds, good seed mixtures without loose peanuts, RSPB food bars and summer seed mixture are all good foods to provide. Soft apples and pears cut in half, bananas and grapes are also good. Some people use soaked dog or cat food and tinned pet foods, but these may attract magpies, crows and cats.

Avoid using peanuts, fat and bread at this time, since these can be harmful if adult birds feed them to their nestlings. If you feel you must put out peanuts, only do so in suitable mesh feeders that will not allow sizeable pieces of peanuts to be removed and provide a choking risk.

Home-made fatballs can go soft and rancid in warm summer weather, and should be avoided. Commercially produced fat bars are suitable for summer feeding but discard any remains after three weeks.

Temporary food shortage can occur at almost any time of the year, and if this happens during the breeding season, extra food on your bird table can make a big difference to the survival of young.

Birds time their breeding period to exploit the availability of natural foods: caterpillars in the case of blackbirds and song thrushes and earthworms in the case of tits and chaffinches. It is now known that if the weather turns cold or wet during spring or summer, severe shortage of insect food can occur, and if the weather is exceptionally dry, earthworms will be unavailable to the ground feeders because of the hard soil.

### **Natural food shortages**

If food shortages occur when birds have young in the nest they may be tempted by easy food put on birdtables to make up the shortfall in natural food, initially to feed themselves, but if the situation gets bad enough, they will also take the food to the nest.

If the food offered on your bird table isn't suitable for the young chicks, it can do more harm than good, and can even be lethal to the chicks as they can choke on the food. It can be difficult for a human to gauge when food shortage in the wild occurs, and hence it is best not to put out food that is likely to create problems during the breeding season.

Therefore, never put out loose peanuts, dry hard foods, large chunks of bread, or fats during the spring or summer months.

© 2009 The Royal Society for the Protection of Birds. Charity registered in England and Wales no 207076, in Scotland no SC037654